

Chord theorems on graphs

Mohammad Javaheri

Department of Mathematics
University of Oregon
Eugene, OR 97403

email: javaheri@uoregon.edu

February 2, 2008

Abstract

The Horizontal Chord Theorem states that if a continuous curve connects points A and B in the plane, then for any integer k there are points C and D on the curve such that $\overrightarrow{AB} = k\overrightarrow{CD}$. In this note, we discuss a few combinatorial-analysis problems related to this theorem and introduce a different formulation that gives way to generalizations on graphs.

1 THE NECKLACE OF PEARLS PROBLEM. *Two pirates have a single-strand necklace containing $2N$ black pearls and $2N$ white pearls arranged in any order. They would like to cut the necklace into as few pieces as possible so that after dividing the pieces of the necklace between them, each gets exactly N white pearls and N black ones.*

Using the following theorem, one can show that two cuts are sufficient. V. Totik has discussed several problems, including the above mentioned and the combinatorial-analysis theorem below, in the Monthly paper ‘*A Tale of Two Integrals*’ [4].

Theorem 1. Suppose f, g are Riemann-integrable functions on $I = [0, 1]$ such that

$$\int_I f(t)dt = \int_I g(t)dt = 1 . \quad (1)$$

Then, for any $r \in [0, 1]$ there exists a subinterval $J \subseteq I$ such that

$$\int_J f(t)dt = \int_J g(t)dt = r \quad \text{or} \quad \int_J f(t)dt = \int_J g(t)dt = 1 - r .$$

According to the arrangement of black and white pearls in the necklace, one defines two step functions $f, g : [0, 1] \rightarrow \mathbb{R}$ such that the equations (1) are satisfied. If J is the interval guaranteed by 1 for $r = 1/2$, then J contains N white pearls and N black pearls (if J happens to have an end point on one of the pearls, then the other end point has to be of the same color and we can modify J accordingly).

Theorem 1 implies that:

Corollary 2. If $f, g : S^1 \rightarrow \mathbb{R}$ are Riemann-integrable functions on the circle such that

$$\int_{S^1} f(\theta)d\theta = \int_{S^1} g(\theta)d\theta = 1 ,$$

then for each $r \in [0, 1]$ there exists an arc U such that

$$\int_U f(\theta)d\theta = \int_U g(\theta)d\theta = r .$$

It is worth mentioning how this corollary is related to the chord set of a function. For a function $F : \mathbb{R} \rightarrow \mathbb{R}$, the chord set of F is defined as

$$H(F) = \{t \in \mathbb{R} : \exists x \in \mathbb{R} \ F(x+t) = F(x)\} .$$

Any $f : S^1 \rightarrow \mathbb{R}$ induces a periodic function p on $\mathbb{R}$ with period one. If $\int_{S^1} f = 0$, then $F(x) = \int_0^x p$ is periodic with period one. Corollary 2 implies that F has every chord, i.e. $H(F) = \mathbb{R}$. In fact, every continuous periodic function has every chord (see [2]).

Another implication of Corollary 2 is that in the following game there is always a winner:

The game board consists of a circle and $2N$ distinguished dots on the circle, $N \geq 1$. The players choose positive integers $m, n \leq N$. The two players

take turns to cross out at least one and at most m dots (that have not been crossed out yet) per turn. One player uses a red pen and the other uses a blue pen (or they use any other method to keep track of the dots they have crossed out). Once a player has crossed out a total number of N dots, the other player has to cross out the remaining dots. The first player who creates an arc of $2n$ consecutive dots with n blue and n red crossed-out dots has lost the game.

The same game can be played on any finite graph. This time players have to avoid creating a *connected subset* of the graph that contains exactly $2n$ dots, n of which are crossed out by the first player and the rest are crossed out by the second player. The game in this general setting on arbitrary graphs may end without a winner. However, if the graph is Euler, then the game always has a winner. A graph is called Euler if it has a closed path that goes through every edge exactly once, or equivalently, it is connected and the degree of every vertex is even. If the graph is Euler, then by considering its Euler circuit, we can pretend the game is played on a circle. It follows from the circle case that the game cannot end without a winner.

2 Partition and Chord sets. Theorem 1 is also related to the Horizontal Chord Theorem which states that if a continuous curve connects points $A, B \in \mathbb{R}^2$, then for each positive integer k there exist points C, D on the curve such that

$$\overrightarrow{CD} = \frac{1}{k} \overrightarrow{AB}.$$

Given functions f and g on I , one considers the plane curve $\gamma(x) = (\int_0^x f, \int_0^x g)$ and concludes from the Horizontal Chord Theorem that there exists an interval J such that $\int_J f = \int_J g = 1/k$ or $\int_J f = \int_J g = -1/k$. A more careful argument (see [1] or [4]) shows that:

Corollary 3. *If $\int_I f(t)dt = \int_I g(t)dt = 1$, then for any positive integer k there exists a subinterval $J \subseteq I$ such that $\int_J f(t)dt = \int_J g(t)dt = 1/k$. Moreover, if k is replaced by a number that is not a positive integer, the conclusion does not necessarily hold.*

Let us now consider the case where $g \equiv 1$ in Corollary 3. By a simple continuity argument, we show that if $\int_I f(t)dt = 1$, then for any positive integer k there exists an interval $J \subset [0, 1]$ of size $1/k$ such that $\int_J f = 1/k$.

To see this, we define

$$F(x) = \int_x^{x+1/k} f(t)dt, \quad x \in [0, 1 - 1/k] .$$

Then

$$F(0) + F\left(\frac{1}{k}\right) + \dots + F\left(1 - \frac{1}{k}\right) = \int_0^1 f(t)dt = 1 .$$

Since F is a continuous function, the Intermediate-value Theorem [3] implies that there exists $x \in [0, 1 - 1/k]$ such that $F(x) = 1/k$, and so $\int_J f = 1/k$ with $J = [x, x + 1/k]$. In this argument, we used a crucial property of $1/k$ with regard to $I = [0, 1]$, namely, there is a partition of $[0, 1]$ to subintervals of size $r = 1/k$. We will generalize this argument in Theorem 5 to all connected finite graphs.

We need to introduce some terminology before presenting the definitions of *partition* and *chord* sets for general graphs. Let $G = (V, E)$ be a connected graph, possibly with loops or multiple edges, where V denotes the set of vertices and E denotes the set of edges of G . We can think of G as a measure space (G, μ_G) by identifying each edge with the interval $[0, 1]$ with the standard uniform measure. The set $L^1(G)$ is the set of functions $f : G \rightarrow \mathbb{R}$ such that f is Riemann-integrable when restricted to each edge.

The identification of each edge in G with the interval $[0, 1]$ also turns G into a topological space. A subset of G is closed if its intersection with each edge of G is closed. Let $X(G)$ denote the set of closed and connected subsets of G . Finally, given $r \in (0, |E|)$, let

$$X_r(G) = \{u \in X(G) : \mu_G(u) = r\} .$$

In section 3, we will see that $X_r(G)$ is a path-connected metric space with the metric:

$$d_G(u, v) = 2r - 2\mu_G(u \cap v), \quad u, v \in X_r(G) .$$

Definition 4. *The partition set of a connected graph G is the set of real numbers r such that there exists $n \in \mathbb{N}$ and a collection $\{U_\alpha\}_1^n \subset X_r(G)$ such that, except for a finite number of points, every point of G appears in exactly n members of the collection. The partition set of G is denoted by $P(G)$.*

The chord set of a connected graph G is the set of real numbers r such that for every $f \in L^1(G)$ with $\int_G f = 0$ there exists $U \in X_r(G)$ such that $\int_U f = 0$. We denote the chord set of G by $H(G)$.

We saw that $P(I) \subseteq H(I)$ for $I = [0, 1]$, the graph with two vertices and one edge. In general:

Theorem 5. *For any finite connected graph G ,*

$$\overline{P(G)} \subseteq H(G) , \quad (2)$$

where $\overline{P(G)}$ is the closure of $P(G)$ in $\mathbb{R}$.

Proof. Since the chord set is a closed subset of $\mathbb{R}$, it is sufficient to prove $P(G) \subseteq H(G)$. Let $r \in P(G)$ and $f \in L^1(G)$ such that $\int_G f = 0$. We will show that there exists $U \in X_r(G)$ such that $\int_U f = 0$.

Let $I_f : X_r(G) \rightarrow \mathbb{R}$ be the integration map defined by

$$I_f(A) = \int_A f d\mu_G .$$

We show that I_f is continuous on $X_r(G)$ (the topology on $X_r(G)$ is induced by the metric d_G). Suppose $A_i \rightarrow A$ in $X_r(G)$. Then

$$|I_f(A_i) - I_f(A)| = \left| \int_{A_i \oplus A} f d\mu_G \right| \rightarrow 0 ,$$

and so I_f is continuous at every $A \in X_r(G)$. Now, since $r \in P(G)$, there exists a collection $\{U_\alpha\}_1^k \subset X_r(G)$ such that almost every point of G appears in exactly n members of the collection. Hence,

$$\sum_{\alpha=1}^k I_f(U_\alpha) = \sum_{\alpha=1}^k \int_{U_\alpha} f d\mu_G = n \int_G f d\mu_G = 0 .$$

The path-connectedness of $X_r(G)$ (see Theorem 7) and the Intermediate-value Theorem [3] imply that there exists $U \in X_r(G)$ such that $I_f(U) = 0$, and (2) follows. $\square$

We conjecture that in fact the equality $\overline{P(G)} = H(G)$ holds.

3 THE SPACE OF CONNECTED SUBSETS. In addition to being a measure space and a topological space, any graph G is equipped with a natural metric. The distance between two points in G is the length of the

shortest path connecting them. We extend this metric on G to $X(G)$ by setting:

$$d_G(A, B) = \mu_G(A \sqcup B) - \mu_G(A \cap B) , \quad A, B \in X(G) ,$$

where $A \sqcup B$ is the smallest closed connected subset of G that contains $A \cup B$. It is straightforward to check that $(X(G), d_G)$ is a metric space. Moreover, $X(G)$ is connected. In fact, any $C \in X(G)$ can be retracted continuously to any $x \in C$, by which we mean:

Lemma 6. *For any nonempty $C \in X(G)$ and any $x \in C$, there exists a continuous curve $\eta : [0, \mu_G(C)] \rightarrow X(G)$ such that $\eta(0) = C$, $\eta(\mu_G(C)) = \{x\}$, $\mu_G(\eta(t)) = \mu_G(C) - t$ for all t , and $\eta(t_1) \subset \eta(t_2)$ whenever $t_1 > t_2$.*

Proof. We prove the assertion by a continuous induction on $\mu(C)$. If $\mu(C) < 1$, then C contains at most one vertex and clearly the retraction of C to any $x \in C$ exists. Suppose the assertion in the lemma is true for all $C \in X(G)$ and all $x \in C$ with $\mu(C) < r$. Let $C \in X(G)$ such that $\mu(C) < r + 1$ and let $x \in C$. If C contains at most one edge, then again the desired retraction clearly exists. Thus, suppose C contains more than one edge. Let a denote the interior of one of the edges in C that does not contain x . Then we can retract C to $C \setminus a$ by removing a continuously from C . By the inductive hypothesis, $C \setminus a$ can be retracted to $x \in C \setminus a$. It follows that C can be retracted continuously to x . $\square$

Our task in this section is to prove the following theorem.

Theorem 7. *The metric space $(X_r(G), d_G)$ is path-connected for $r \in (0, |E|)$.*

Proof. Let $A, B \in X_r(G)$ and let $\mathcal{C}$ be the path-connected component of $X_r(G)$ that contains A . We first show that there exists an element in $\mathcal{C}$ that intersects with B . Let

$$\alpha = \inf_{U \in \mathcal{C}} d_G(B, U) . \quad (3)$$

Choose $C \in X_r(G)$ such that

$$d_G(B, C) < \alpha + r . \quad (4)$$

If $B \cap C = \emptyset$, then there exists a continuous curve $\gamma : [0, s] \rightarrow G$ with $\mu_G(\gamma([0, t])) = t$, for all $t \in [0, s]$, such that $\gamma(0) \in C$ and $\gamma(s) \in B$, where $s = d_G(B, C) - 2r > 0$. Let

$$\theta = \min\{r, s\} .$$

Let C_t , $t \in [0, r]$, be the retraction of C to $\gamma(0) \in C$, obtained in Lemma 6. Then

$$U_t = C_t \cup \gamma([0, t]) \quad , \quad t \in [0, \theta]$$

is a well-defined continuous curve in $X_r(G)$ that connects C to C_θ . Now, we have

$$d_G(B, C_\theta) \leq d_G(B, C) - \theta = s + 2r - \theta . \quad (5)$$

If $s > \theta$, then $\theta = r$, and the inequalities (4) and (5) imply that $d_G(B, C_\theta) < \alpha$ which contradicts (3). Hence, $s = \theta$ and $d_G(B, C_\theta) = 2r$, i.e. $B \cap C_\theta \neq \emptyset$.

Nex, we fix $x \in B$ such that $\mathcal{C}_x = \{C \in \mathcal{C} : x \in B \cap C\} \neq \emptyset$. By the discussion above, we know such x exists. For each $C \in \mathcal{C}_x$, let U_C be the connected component of $B \cap C$ that contains x and set

$$T = C \setminus U_C = \bigcup_{i=1}^k T_i \quad , \quad B \setminus C = \bigcup_{i=1}^l B_i \quad ,$$

where each B_i (respectively, T_i) is a connected component of $B \setminus C$ (respectively, T). The numbers $k = k(C)$ and $l = l(C)$ are functions of C . Define

$$\omega = \omega(C) = k(C) + l(C) \geq 0 .$$

We show that if $\omega(C) > 0$, then there exists $C' \in \mathcal{C}_x$ such that $\omega(C') \leq \omega(C) - 1$. Without loss of generality, we can assume there exists $y \in \overline{B_1} \cap U_C$. Also there exists $z \in \overline{T_1} \cap U_C$, since C is connected. Let $\eta : [0, \mu_G(T_1)] \rightarrow G$ be the retraction of T_1 to z obtained in Lemma 6. Similarly, let $\nu : [0, \mu_G(B_1)] \rightarrow G$ be the retraction of B_1 to y . Let

$$V_t = (C \setminus (T_1 \setminus \eta(t))) \cup \nu(\mu_G(B_1) - t) \quad , \quad t \leq \min\{\mu_G(B_1), \mu_G(T_1)\} .$$

Then V_t is a continuous curve in $X_r(G)$ connecting C to $C' = V_\lambda$, where $\lambda = \min\{\mu_G(B_1), \mu_G(T_1)\}$. If $\lambda = \mu_G(B_1)$, then $l(C') \leq l(C) - 1$. On the other hand, if $\lambda = \mu_G(T_1)$ then $k(C') \leq k(C) - 1$. In either case:

$$\omega(C') = k(C') + l(C') \leq \omega(C) - 1 \quad , \quad (6)$$

Now replace C by C' and repeat the process above. In a finite number of steps we get $C' \in \mathcal{C}$ with $\omega(C') < 1$. But then $k(C') = 0$ and so $B = C' \in \mathcal{C}$. This concludes the proof of Theorem 7. $\square$

4 A MORE GENERAL RESULT. Corollary 2 implies that if $G = (V, E)$ is Euler, then G has every chord, i.e. $H(G) = [0, |E|]$. In this section, we prove a result that holds on more general graphs:

Theorem 8. *Suppose G is a finite connected graph (possibly with loops and parallel edges) with no vertex of degree 1. Then*

$$[0, 1] \subseteq H(G) .$$

A closed path is called a *semi-simple* closed path if i) no two consecutive edges along the path are the same and ii) every edge is repeated at most twice along the path. We say a graph G admits a double covering by semi-simple closed paths, if there are semi-simple closed paths C_i , $i = 1, \dots, k$, that altogether cover every edge of G exactly twice. In other words, every edge in G either belongs to exactly two C_i 's but is repeated once in each one of them, or it belongs to only one of the C_i 's but is repeated twice. We allow repetitions among C_i 's, i.e. it is possible that C_i and C_j contain the same edges and $i \neq j$.

It is easily seen that if G contains a vertex of degree one, then G does not admit a double covering by semi-simple closed paths. In order to prove Theorem 8, we need to show that the converse is true:

Lemma 9. *Suppose G is a connected graph such that*

$$\deg v > 1, \forall v \in V(G) . \tag{7}$$

Then G admits a double covering by semi-simple closed paths.

Proof. Let $\mathcal{S}$ denote the set of all subgraphs of G that admit a double covering by semi-simple closed paths. The degree condition (7) implies that there exists a simple closed path in G . Since every simple closed path is a semi-simple closed path, the set $\mathcal{S}$ is nonempty. Let K be a maximal element of $\mathcal{S}$, i.e. K is not a subgraph of any other element of $\mathcal{S}$. Let L be the graph obtained from G by removing the edges of K . Since K is maximal in $\mathcal{S}$, the subgraph L does not contain any closed paths. In particular, there are no loops or parallel edges in L . In the sequel, we prove by contradiction that L contains no edges.

Let α be the longest simple path in L and let u and v be the endpoints of α . It follows from the degree condition (7) and the fact that there are no cycles in L that u and v are distinct vertices of K . Since $K \in \mathcal{S}$, there

is a collection $M = \{C_i\}_1^k$ of semi-simple closed paths such that every edge of G appears altogether twice along these paths. In particular, there are semi-simple closed paths $\gamma = C_i$ and $\eta = C_j$ that contain the vertices u and v respectively. If $i \neq j$, then the path $\beta = \gamma \cdot \alpha \cdot \eta \cdot \overleftarrow{\alpha}$ is a semi-simple path, where $\overleftarrow{\alpha}$ means the reverse of α . We can remove γ and η from the collection M and add β to obtain a larger element in $\mathcal{S}$. This contradicts our assumption that K was maximal. If $i = j$, then we partition γ to two paths γ_1 and γ_2 such that γ_1, γ_2 are paths from u to v . Now by removing γ from the collection M and adding the path $\gamma_1 \cdot \alpha \cdot \gamma_2 \cdot \alpha$ we obtain a larger element of $\mathcal{S}$. This again contradicts the maximality of K . It follows that L contains no edges, i.e. $K = G$ and G admits a double covering by semi-simple closed paths. $\square$

For a path $P = a_1 \dots a_m$ and $f \in L^1(G)$, we define

$$\int_P f d\mu_G = \sum_{i=1}^m \int_{a_i} f d\mu_G .$$

Now we are ready to present the proof of Theorem 8.

Proof of Theorem 8. Let $r \in [0, 1]$ and $f \in L^1(G)$ such that $\int_G f = 0$. By Lemma 9 and by the definition of double covering, there exist semi-simple closed paths $C_1, \dots, C_k$ such that:

$$\sum_{i=1}^k \int_{C_i} f d\mu_G = 2 \int_G f d\mu_G = 0 \quad (8)$$

We note that if C is a semi-simple closed path, then there exist $A \in X_r(G)$ such that $\int_A f = r \int_C f$. To see this, suppose the length of C is n and let F_n denote the cycle of length n . Then there is a continuous map $\phi : F_n \rightarrow C$ that maps edges of F_n to edges of C . Moreover, we can choose ϕ to be measure-preserving when restricted to each edge of F_n . By Corollary 2, there is $B \in X_r(F_n)$ such that

$$\int_B f \circ \phi(\theta) d\theta = r \int_{F_n} f \circ \phi(\theta) d\theta . \quad (9)$$

Since $r \leq 1$ and C is semi-simple, the set $A = \theta(B)$ is connected and has size r . Moreover, we conclude from (9) that $\int_A f = \int_B f \circ \phi = r \int_C f$.

It follows that for each $i = 1, \dots, k$ there exists $A_i \in X_r(G)$ such that

$$I_f(A_i) = \int_{A_i} f d\mu_G = r \int_{C_i} f d\mu_G . \quad (10)$$

The equations (8) and (10) imply that

$$\sum_{i=1}^k I_f(A_i) = r \sum_{i=1}^k \int_{C_i} f d\mu_G = 0 .$$

Then the Intermediate-value Theorem implies that there exist $U \in X_r(G)$ such that $I_f(U) = 0$ and so $r \in H(G)$. $\square$

References

- [1] *Contests in Higher Mathematics, 1949-1961*, Akadémiai Kiadó, Budapest, 1968.
- [2] J.C. Oxtoby, *Horizontal Chord Theorem*, Amer. Math. Monthly **79** (1972) 468–475.
- [3] K.A. Ross, *Elementary Analysis: The Theory of Calculus*, Springer, 1980.
- [4] V. Totik, *A Tale of Two Integrals*, Amer. Math. Monthly **106** (1999) 227–240.